

Global Assessment of Land Degradation and Improvement

1. Identification by remote sensing

Z G Bai
D L Dent
L Olsson
M E Schaepman



World Soil Information



FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS

This report has been prepared under the conditions laid down in the Letter of Agreement
FAO-ISRIC PR 35852

Citation:

Bai ZG, Dent DL, Olsson L and Schaepman ME 2008. *Global assessment of land degradation and improvement. 1. Identification by remote sensing.* Report 2008/01, ISRIC – World Soil Information, Wageningen

Inquiries:

C/o Director, ISRIC – World Soil Information
PO Box 353
6700 AJ Wageningen
The Netherlands
Telefax: +31 (0)317 471700
E-Mail: soil.isric@wur.nl
Web: www.isric.org

*I have been a field for nigh on a thousand years, and I know men.
Some are clever, some are kind, but very few are clever and kind, but he was.
And I am sorry that all the other fields of England – who need him so much these
days - will have to go on without him.*

Obituary for a Suffolk farmer 1930

PRELIMINARY FINDINGS

Land degradation is a global environment and development issue. Up-to-date, quantitative information is needed to support policy and action for food and water security, economic development, environmental integrity and resource conservation. To meet this need, the Global Assessment of Land Degradation and Improvement (GLADA) uses remote sensing to identify degrading areas and areas where degradation has been arrested or reversed. This screening will be followed up within the parent LADA program by field investigations to establish the situation on the ground. Findings remain provisional until validated in the field.

1. **Land degradation is defined as a long-term decline in ecosystem function and measured in terms of net primary productivity.** The remotely-sensed normalized difference vegetation index (NDVI) is used as a proxy; its deviation from the norm may serve as an indicator of land degradation and improvement if other factors that may be responsible (climate, soil, terrain and land use) are accounted for. Rainfall effects may be accounted for by rain-use efficiency (NDVI per unit of rainfall) and residual trends of NDVI, temperature effects by energy-use efficiency (derived from annual accumulated temperature). Land degradation is indicated by a declining trend of climate-adjusted net primary productivity and land improvement by an increasing trend. Translation of NDVI to net primary productivity enables economic appraisal of land degradation.
2. **Land degradation is cumulative – this is the global issue.** The 1991 GLASOD assessment indicated that 15 per cent of the land surface was degraded; the present assessment identifies 24 per cent of land as degrading but the areas hardly overlap, which means that new areas are being affected. Some areas of historical land degradation have been so degraded that they are now stable - at stubbornly low levels of productivity.
3. **Analysis of 23-year GIMMS NDVI data reveals a declining trend across some 24 per cent of the global land area.** Spatial patterns and temporal trends of NDVI and rain-use efficiency are analysed for the period 1981-2003 at 8km resolution. Degrading areas are mainly in Africa south of the Equator, SE Asia and S China, N-Central Australia, the Pampas, and swaths of boreal forest in Siberian and N America.
4. **Almost one fifth of degrading land is cropland - more than 20 per cent of all cultivated areas;** 23 per cent is broadleaved forest, 19 per cent needle-leaved forests, 20-25 per cent rangeland. Cropland occupies only 12 per cent of the land area and forest only 28 per cent, so degradation is over-represented in both cropland and in forest globally.
5. **Loss of carbon fixation from the atmosphere, associated with land degradation over the period, amounts to almost a thousand million tonnes.** At a shadow price of \$50 per tonne, the cost is almost \$50 billion. The cost of land degradation is at least an order of magnitude greater in terms of *emissions* to the atmosphere though loss of soil organic carbon.

6. **Some 16 per cent of the land area shows improvement.** 18 per cent of the improving area is cropland (20 per cent of the total croplands), 23 per cent is forest and 43 per cent rangeland.
7. **There is only a weak correlation with biophysical factors other than land cover:** 78 per cent of degrading land is in humid regions, 8 per cent in the dry sub-humid, 9 per cent in the semi-arid, and 5 per cent in arid and hyper-arid regions. There is no obvious relationship between degrading land and the nature of soil or terrain – degradation is driven by management and catastrophic natural phenomena.
8. **About 1.5 billion people depend directly on the degrading areas.** There is a weak correlation between degrading land and rural population density but more detailed analysis of land use history is needed to tease out the underlying social and economic drivers.

Key words: land degradation/improvement, remote sensing, NDVI, net primary productivity, land use/cover, global relationships

Contents

PRELIMINARY FINDINGS	i
Abbreviations	v
1 Introduction	1
1.1 The need for a new assessment.....	1
1.2 Indicators.....	1
2 Data and methods	5
2.1 Data.....	5
2.2 Analysis	7
3 Results	9
3.1 Greenness trends	9
3.2 Climate-adjusted greenness.....	11
3.2.1 Rain-use efficiency	11
3.2.2 RESTREND	17
3.2.3 Energy-use efficiency.....	17
4 Land degradation and improvement	23
4.1 Land degradation	23
4.1.1 Which regions are hardest hit?	23
4.1.2 Is land degradation a global issue?	28
4.1.3 Is land degradation mainly associated with farming?	29
4.1.4 Is land degradation a dryland issue?	35
4.1.5 Is it related to population pressure?.....	35
4.1.6 Is land degradation related to poverty?	35
4.2 Land improvement	36
4.3 Urban areas.....	36
4.4 Comparison of indicators	36
5 What GLADA can and cannot show; further analysis	39
6 Conclusions	43
Acknowledgements	44
References	45
Appendix 1: Data	51
Appendix 2: Analytical methods	53
Appendix 3: NDVI indicators of the land degradation / improvement ..	55

Figures

Figure 1.	Spatially aggregated annual sum NDVI 1981-2003, $p < 0.01$	9
Figure 2.	Global change in net primary productivity, 1981-2003.....	10
Figure 3.	Global change in rain-use efficiency, 1981-2003.....	12
Figure 4.	Correlation between annual sum NDVI and annual rainfall, 1981-2003	13
Figure 5.	Global negative trend in RUE-adjusted NDVI, 1981-2003.....	15
Figure 6.	NPP loss in the degrading areas 1981-2003	16
Figure 7.	Residual trends of sum NDVI (RESTREND), 1981-2003.....	18
Figure 8.	Confidence levels of RESTREND, 1981-2003	19
Figure 9.	Global change in energy-use efficiency, 1981-2002.....	20
Figure 10.	Global trend of positive climate-adjusted NDVI, 1981-2003.....	21
Figure 11.	Confidence levels of positive climate-adjusted NDVI, 1981-2003.....	22
Figure 12.	South Africa, land degradation and population affected, 1981-2003 ...	34
Figure 13.	South Africa: relationship between population density and land degradation / improvement.....	35
Figure 14.	Eastern South Africa: sum NDVI, RUE-adjusted NDVI, rain-use efficiency, and RESTREND	38

Tables

Table 1:	Statistics of degrading areas 1981-2003, by country.....	24
Table 2.	Land degradation in LADA partner countries by global rank order.....	28
Table 3.	Global degrading and improving areas by land cover.....	30
Table 4.	Global degrading and improving areas by land use systems	31
Table 5.	Global degrading/improving lands in the aggregated land use systems	33
Table 6.	Global comparison of trends in various indicators	37

Abbreviations

AVHRR	Advanced Very High Resolution Radiometer
Biome-BGC	Biome-BioGeochemical Cycles, Terrestrial Ecosystem Process Model
CIAT	International Centre for Tropical Agriculture, Cali, Columbia
CIESIN	Center for International Earth Science Information Network, Colombia University, Palisades, NY
CoV	Coefficient of Variation
CRU TS	Climate Research Unit, University of East Anglia, Time Series
ENSO	El Niño/Southern Oscillation
EUE	Energy-Use Efficiency
FAO	Food and Agriculture Organization of the United Nations, Rome
GEF	The Global Environment Facility, Washington DC
GIMMS	Global Inventory Modelling and Mapping Studies, University of Maryland
GLADA	Global Assessment of Land Degradation and Improvement
GLASOD	Global Assessment of Human-Induced Soil Degradation
GPCC	The Global Precipitation Climatology Centre, German Meteorological Service, Offenbach
HANTS	Harmonic Analyses of NDVI Time-Series
JRC	European Commission Joint Research Centre, Ispra, Italy
LADA	Land Degradation Assessment in Drylands
Landsat ETM+	Land Resources Satellite, Enhanced Thematic Mapper
LUS	Land Use Systems, FAO
MOD17A3	MODIS 8-Day Net Primary Productivity data set
MODIS	Moderate-Resolution Imaging Spectroradiometer
NDVI	Normalized Difference Vegetation Index
NPP	Net Primary Productivity
RESTREND	Residual Trends of sum NDVI
RUE	Rain-Use Efficiency
SOTER	Soil and Terrain database
SPOT	Système Pour l'Observation de la Terre
SRTM	Shuttle Radar Topography Mission
TRMM	The Tropical Rainfall Measuring Mission
UNCED	United Nations Conference on Environment and Development
UNEP	United Nations Environment Programme, Nairobi
VASCLimO	Variability Analyses of Surface Climate Observations

1 Introduction

1.1 The need for a new assessment

Economic development, burgeoning cities and increasing rural populations, are driving unprecedented land-use change. In turn, unsustainable land use is driving land degradation – a long-term loss in ecosystem function and productivity which requires progressively greater inputs to recoup the situation. Its symptoms include soil erosion, nutrient depletion, salinity, water scarcity, pollution, disruption of biological cycles, and loss of biodiversity. This is a global development and environment issue recognised by the UN Convention to Combat Desertification, the Conventions on Biodiversity and Climatic Change, and the Millennium Goals (UNCED 1992, UNEP 2007).

Quantitative, up-to-date information is needed to support policy development for food and water security, environmental integrity, and economic development. But land degradation is a contentious field. Crucial questions that must be answered in a scientifically justifiable way include: Is land degradation a global issue or a collection of local problems? Which regions are hardest hit; how hard are they hit? Is it mainly a problem of drylands? Is it mainly associated with farming? Is it related to population pressure - or poverty? A globally consistent assessment also serves as a yardstick for measuring the seriousness of any local situation.

The only previous harmonized assessment, the *Global Assessment of Human-induced Soil Degradation (GLASOD)*, was a compilation of expert judgements of the kind and degree of land degradation, e.g. soil erosion by water or by wind, salinity, nutrient depletion (Oldeman and others 1991). It was a map of perceptions, not a measure, of land degradation and is now out-of-date; its qualitative judgments (Appendix Table S1) have proven inconsistent and hardly reproducible, relationships between land degradation and policy-pertinent criteria were unverified (Sonneveld and Dent 2007) - as its authors were the first to point out. Within the FAO program *Land Degradation Assessment in Drylands (LADA)*, the present global assessment uses remote sensing to identify areas where significant biological change is happening, both *hot spots* of land degradation and *bright spots* of land improvement. In the next phase of the program, hot spots and bright spots will be further characterised in the field by national teams; findings remain provisional until validated in the field.

1.2 Indicators

Land degradation may be defined as a long-term loss of ecosystem function and productivity caused by disturbances from which the land cannot recover unaided. It may be measured by change in net primary productivity (NPP - the rate at which vegetation fixes CO₂ from the atmosphere less losses through respiration); deviation from the norm may be taken as an indicator of land degradation or

improvement. As a proxy, the remotely sensed normalized difference vegetation index (NDVI) has been shown to be related to biophysical variables that control vegetation productivity and land/atmosphere fluxes (Hall and others 2006) such as: leaf-area index (Myneni and others 1997), the fraction of photosynthetically-active radiation absorbed by vegetation (Asrar and others 1984), and NPP (Alexandrov & Oikawa 1997, Rasmussen 1998a, b). It has also been used to estimate vegetation change, either as an index (Anyamba & Tucker 2005, Olsson and others 2005) or as one input to dynamic vegetation models (Nemani and others 2003, Seaquist and others 2003, Fensholt and others 2006). Consistent time-series data at spatial resolutions from 20m to 8km (Brown and others 2006) enable analysis and generalization.

This study uses NDVI data produced by the Global Inventory Modelling and Mapping Studies (GIMMS) group from measurements made by the AVHRR radiometer on board US National Oceanic and Atmospheric Administration satellites. The fortnightly images at 8km spatial resolution are corrected for calibration, view geometry, volcanic aerosols, and other effects not related to vegetation cover (Tucker and others 2004).

A negative trend in NDVI does not necessarily indicate land degradation, nor does a positive trend necessarily indicate land improvement. Biomass depends on several factors including: climate - especially fluctuations in rainfall, sunshine, and length of growing season; land use; large-scale ecosystem disturbances such as fires; and the global increase in nitrate deposition and atmospheric carbon dioxide. To interpret NDVI trends in terms of land degradation or improvement, we have to eliminate false alarms, in particular those arising from climatic variability and land use change. Globally, this can be done for climate, for which a century's consistent data are available, but global time series are not available for land use which has to be examined case-by-case.

Where productivity is limited by rainfall, rain-use efficiency (RUE, the ratio of NPP to rainfall) accounts for variability of rainfall and, to some extent, local soil characteristics (Houérou 1984, Houérou and others 1988). The combination of satellite-based estimation of NDVI and station-observed rainfall has been used to assess land degradation at various scales (Holm and others 2003, Prince and others 2007) but RUE, itself, is strongly correlated with rainfall. In the short term, it says more about rainfall fluctuation than about land degradation but we judge that its long-term trends distinguish between the effects of rainfall variability and land degradation. To get around the correlation of RUE with rainfall, Wessels and others (2007) have suggested the alternative use of residual trends – the difference between the observed NDVI and that predicted from the local rainfall-NDVI relationship. Both approaches are employed in this report.

There are caveats when applying these data globally:

- 1) The NDVI signal can be saturated at closed vegetation canopy (Ripple 1985). This means that NDVI is more sensitive for cropland and rangeland than for forest; however, it is still useful for forest;
- 2) Cloud screening was performed on the GIMMS data and maximum NDVI was read out for a composite of 15 days, but NDVI may still be underestimated for very cloudy areas;

- 3) The great spatial variability of rainfall in drylands makes interpolation of point measurements problematic, and observation stations are sparse in many of these areas.

NDVI is simply a ratio of red and near-infrared light reflected by the land surface. To provide a more tangible measure of land degradation that may be subjected to economic analysis, the GIMMS data are translated to NPP using MODIS (moderate-resolution imaging spectroradiometer) NPP data (Running and others 2004) for the overlapping period 2000-2003; this translation is approximate. From the year 2000, NPP has been calculated from MODIS measurements of the fraction of photosynthetically-active radiation absorbed by vegetation (which does not saturate at high leaf areas) at 1km resolution; this is the preferred indicator for the future.

NDVI cannot be other than a proxy for land degradation and improvement. We are using this indicator simply to identify areas where significant biological change is taking place; land degradation is identified by a declining trend in climate-adjusted NDVI and land improvement by a rising trend. It does not tell us anything about the nature of the changes; what is happening in, say, south China is different from what is happening in the Pampas, both in terms of the driving changes in land use and the kind of land degradation. The patterns of land degradation and improvement, so identified, are further explored by comparisons with land cover, soil and terrain, and socio-economic data. In the parent LADA program, areas identified in this screening will be validated and characterized in the field by national teams.

2 Data and methods

2.1 Data

GIMMS (Global Inventory Modelling and Mapping Studies) radiometer (AVHRR) data are collected by National Oceanic and Atmospheric Administration satellites. They are corrected for calibration, variations in solar and view zenith angle, El Chichon and Mt Pinatubo stratospheric aerosols, and other effects not related to vegetation change, and generalized to 8km grids for 15-day periods. Global data are currently available for the period July 1981-December 2003 (Tucker and others 2004).

NDVI-NPP correlation: To get a measure open to economic analysis, the GIMMS NDVI time series has been translated to NPP using MODIS data (Justice and others 2002, Running and others 2004)¹ for the overlapping period 2000-2003. NPP was estimated by correlation with MODIS 8-day NPP values for the overlapping years of the GIMMS and MODIS datasets (2000-2003), re-sampling the annual mean MODIS NPP at 1km resolution to 8km resolution using nearest-neighbour assignment. The empirical relationship is:

$$\text{NPP}_{\text{MOD17}} [\text{kgC ha}^{-1} \text{ year}^{-1}] = 1106.37 * \text{sum NDVI} - 564.55 \quad [1]$$

($r = 0.83$, $n = 3\,128\,207$)

Where $\text{NPP}_{\text{MOD17}}$ is annual mean NPP derived from MODIS MOD17 Collection 4 data, and sum NDVI is the four-year (2000-2003) mean annual sum NDVI derived from GIMMS. Uncertainty is for slope ± 3.818 , and for intercept ± 16.364 .

VASclimO 1.1 comprises the most complete monthly precipitation data for 1951-2000, compiled from long, quality-controlled station records, gridded at resolution of 0.5° , from 9 343 stations (Beck and others 2005). For 2001-2003, these were supplemented by the GPCP full data re-analysis product (Schneider and others 2008) to produce monthly rainfall values matching the GIMMS NDVI data.

CRU TS 2.1 comprises monthly values of station-observed meteorological data from the beginning of the 20th century, gridded at 0.5° resolution (Mitchell and Jones 2005). Monthly temperature values since January 1981 were used for this analysis.

Rain-use efficiency (RUE), represented by the ratio of annual sum NDVI and annual rainfall, was calculated from the VASclimO rainfall data.

¹ MOD17A3 is a dataset of terrestrial gross and net primary productivity computed at 1-km resolution and an 8-day interval. Though far from perfect (Plummer 2006), MODIS gross and net primary productivity values are related to observed atmospheric CO₂ and the inter-annual variability associated with the ENSO phenomenon, indicating that the NPP data are reliable at the regional scale (Zhao and others 2005, 2006). The dataset has been validated in various landscapes (Fensholt and others 2004, 2006, Gebremichael and Barros 2006, Turner and others 2003, 2006).

Energy-use efficiency (EUE), represented by the ratio of annual sum NDVI to annual accumulated temperature (day degrees above 0°C), calculated from CRU 2.1 monthly data.

Trends analysis: Trends of NDVI and its derivatives were determined by linear regression; the absolute change (Δ) is the slope of the regression. The data were tested for temporal and spatial independence following Livezy and Chen (1983): when the absolute values of the autocorrelation coefficients of lag-1 to lag-3, calculated for a time series consisting of n observations, are not larger than the typical critical value, i.e. $1.96/\sqrt{n}$ corresponding to 5 per cent significance level, the observations in this time series may be accepted as being independent from each other. The T- test was used to arrange the slope values in classes showing strong or weak positive or negative trends:

$$T = b / se(b)$$

Where b is the estimated slope of the regression line between the observation values and time and $se(b)$ represents the standard error of b .

The class boundaries were defined for 99, 95 and 90 per cent confidence levels.

RESTREND: Following the general procedure of Wessels and others (2007), correlation between annual sum NDVI and annual rainfall was calculated for each pixel. For the northern hemisphere, annual sum NDVI was calculated for the calendar year, for the southern hemisphere beginning October 1 through the following September. The regression equation enables prediction of sum NDVI according to rainfall. Residuals of sum NDVI (i.e. differences between the observed and predicted sum NDVI) for each pixel were calculated, and the trend of these residuals (RESTREND) was analysed by linear regression.

Aridity index was calculated as P/PET where P is annual precipitation in mm and $PET = P / \sqrt{(0.9 + (P/L)^2)}$ where $L = 300 + 25T + 0.05T^3$ and T is mean annual temperature (Jones 1997). Precipitation data were taken from the VASCLimO dataset and data for mean annual temperature from the CRU dataset.

Soil and terrain: For this study, a global soil and terrain database at scale 1:1 million-scale has been compiled using the 90m-resolution SRTM digital elevation model and dataset of key soil attributes (SOTER) for the LADA partner countries has been prepared (ISRIC 2008a, b). Correlations between land degradation and soil and terrain have so far been investigated only for Kenya, Cuba and South Africa - using existing SOTER databases.

Land cover: GLC2000 global land cover data (JRC 2003) have been generalised for preliminary comparison with NPP trends.

Land use systems: Land use systems of the World (FAO 2008) have been used for preliminary comparison with NPP trends.

Population, urban areas and poverty indices: The CIESIN Global Rural-Urban Mapping Project provides data for population and urban extent, gridded at 30 arc-second resolution (CIESIN 2004). Sub-national rates of infant mortality and child underweight status and the gridded population for 2005 at 2.5 arc-minutes resolution (CIESIN 2007) were compared with indices of land degradation.

Comparisons between land degradation and other indices: Maps of the climate-adjusted NDVI index were overlaid on the other global maps. Corresponding comparative values were calculated, and correlation calculated for all pixels.

2.2 Analysis

Areas of land degradation and improvement are identified by a sequence of analyses of the remotely sensed data:

1. Simple NDVI indicators (NDVI minimum, maximum, maximum-minimum, mean, sum, standard deviation and coefficient of variation) are computed for the calendar year for the northern hemisphere, and for October to the following September for the southern hemisphere, encompassing a complete growing season. Each of these indicators has biological meaning (Appendix 3).
2. The annual sum NDVI, the aggregate of greenness over the growing season, is used as the standard surrogate for annual biomass productivity. NDVI is translated to net primary productivity by correlation with MODIS data; trends are calculated by linear regression.
3. To distinguish between declining productivity caused by land degradation, and declining productivity due to other factors, false alarms must be eliminated. Rainfall variability and irrigation have been accounted for by:
 - a. Identifying each pixel where there is a positive relationship between productivity and rainfall, i.e. where rainfall determines NPP;
 - b. For those areas where rainfall determines NPP, RUE has been considered: where productivity declined but RUE increased, we may attribute declining productivity to declining rainfall; those areas are masked (urban areas are also masked);
 - c. For the remaining areas with a positive relationship but declining RUE, and also for all areas where there is a negative relationship between NDVI and rainfall, i.e. where rainfall does not determine NPP, NDVI trend has been calculated; this is called *RUE-adjusted NDVI*;
 - d. Land degradation is indicated by a negative trend of RUE-adjusted NDVI and may be quantified as *RUE-adjusted NPP*.
4. As an additional indicator, the residual trend of sum NDVI (RESTREND) is calculated.

5. To take account of the significant lengthening and warming of the growing season at high latitudes and high altitudes, energy-use efficiency (EUE) is calculated for all pixels. In practice, it scarcely affects the estimation of land degradation but does affect the identification of land improvement which is indicated by a positive trend in both RUE-adjusted NPP and EUE, and is quantified as *climate-adjusted NPP*.
6. The indices of land degradation and improvement are compared with land cover; soil and terrain; rural population density; and indices of aridity and poverty.

Algorithms have been devised to undertake these screening analyses automatically. Details of the analytical methods are given as Appendix 2.

3 Results

3.1 Greenness trends

Globally over for the period 1981-2003, greenness increased by 3.8 per cent ($P < 0.05$) but there are significant variations at the continental scale (Figure 1) and at the country scales (Bai and Dent 2007 a-f). The increase was 3 per cent in Africa and North America, 4.4 per cent in Latin America, 4.5 per cent in Australia, 5.4 per cent in Europe, and 6 per cent in Asia. Several regional patterns track the ENSO cycle - with losses during El Niño events and gains during La Nina events. Figure 2 depicts global change in NDVI, scaled in terms of NPP, over the period 1981-2003; ice and extreme desert with NPP less than 1gC m^{-2} are designated as *no change*.




Figure 1. Spatially aggregated annual sum NDVI 1981-2003, $p < 0.01$




Figure 2. Global change in net primary productivity, 1981-2003

3.2 Climate-adjusted greenness

Rainfall variability is taken into account in two ways: by adjustment of NDVI/NPP for rain-use efficiency, and by calculation of residual trends of departures of observed NDVI from values predicted by the rainfall-NDVI relationship (RESTREND). In addition, NDVI/NPP was adjusted for energy-use efficiency using global temperature data.

3.2.1 Rain-use efficiency

Rain-use efficiency (RUE) is production per unit of rainfall. It may fluctuate dramatically in the short term; often there is a sharp decline in RUE when rainfall increases and we assume that the vegetation, whether cultivated or semi-natural, cannot make immediate use of the additional rain. But where rainfall is the main limiting factor on biomass productivity, we judge that the long-term trend of RUE is a good indicator of land degradation or improvement (Houérou 1984, Houérou and others 1988, Snyman 1998, Illius and O'Connor 1999, O'Connor and others 2001). Furthermore, pixel-by-pixel analysis of RUE accommodates the effects of local variations in terrain, soil and vegetation (Justice and others 1991).

In North China and Kenya, Bai and others (2005, 2006) demonstrated that RUE values calculated from NDVI, which are easy to obtain, were comparable with those calculated from field measurements of NPP, which are not easy to obtain. Globally, RUE was calculated as the ratio between annual sum NDVI and station-observed annual rainfall. Figure 3 maps global trends of RUE over the period 1981-2003.

Figure 4 depicts relationship of sum NDVI with rainfall. Drylands mostly show a positive relationship between RUE and rainfall, i.e. biomass production is determined by the amount of rainfall; humid and cold regions, irrigated areas and some wetlands mostly show a negative relationship, i.e. biomass production is not limited by rainfall; but there are some exceptions that may be related to land use change and/or land degradation.




Figure 3. Global change in rain-use efficiency, 1981-2003




Figure 4. Correlation between annual sum NDVI and annual rainfall, 1981-2003

RUE was used to adjust NDVI/NPP values as follows:

- a. For those pixels where there is a positive relationship (i.e. where production is limited by rainfall), where productivity declined but RUE increased, we attribute the decline in productivity to declining rainfall; those areas were masked;
- b. For the remaining areas with a positive relationship *but declining RUE* and, also, pixels where there is a negative relationship between NDVI and rainfall (taken to be areas of rainfall surplus compared with transpiration needs, or irrigated, or areas depending on groundwater), NDVI trends were calculated as *RUE-adjusted NDVI*.

As a first cut, we may equate declining RUE-adjusted NDVI with land degradation (Figure 5a). The confidence levels of these negative trends in RUE-adjusted NDVI shows that two per cent of the land area exhibits a negative trend at the 99% confidence level, 5 per cent at 95% confidence and 7.5 % at the 90% confidence level. The smallness of these areas may be explained by the coarse resolution of the GIMMS data (8km): an area of land degradation much smaller than 8km across must be severe indeed to be seen through the signal from a much larger surrounding area; we see through a glass darkly.

These results have been validated by field observation in North China (Bai and others 2005) and independently by Chen & Rao (2008); Kenya (Bai & Dent 2006); and Bangladesh (Bai 2006).

Figure 6 shows a pixel-based estimate of the loss of NPP compared with the average over the period 1981-2003.




Figure 5. Global negative trend in RUE-adjusted NDVI, 1981-2003




Figure 6. NPP loss in the degrading areas 1981-2003

3.2.2 **RESTREND**

RUE is strongly correlated with rainfall and fluctuates over short periods along with fluctuations of rainfall. To get around these relationships, Wessels and others (2007) suggest the alternative use of Residual Trends to distinguish land degradation from the effects of rainfall variability.

Following their general procedure, we correlated annual sum NDVI and annual rainfall for each pixel (Figure 4); the resulting regression equation allows prediction of sum NDVI according to rainfall in any one year. Residuals of sum NDVI (differences between the observed and predicted sum NDVI) were calculated for each pixel and residual trend (RESTREND) was analysed by linear regression (Figure 7); its significance was assessed by the T-test (Figure 8).

RESTREND points in the same direction as RUE: negative values may indicate human-induced land degradation and positive values, improvement (cf Section 4.4).

3.2.3 **Energy-use efficiency**

Energy use efficiency (EUE) is calculated as the ratio of annual sum NDVI to accumulated temperature (day-degrees Celsius above zero). Figure 9 shows its trend over the period 1981-2002. The global increase in temperatures, especially at high latitudes, has been accompanied by a marked increase in NDVI (Figure 2) but not, in general, in the EUE of either natural vegetation or farmland.

Combination of negative EUE indicator with negative RUE-adjusted NDVI makes virtually no difference to the delineation of land degradation – areas of negative EUE are also areas of negative RUE-adjusted NDVI. However, addition of the EUE indicator does make a big difference to the assessment of land improvement: Figure 10 maps the areas that exhibit *both* a positive trend in RUE-adjusted NDVI and positive EUE as climate-adjusted NDVI; Figure 11 depicts the confidence levels.




Figure 7. Residual trends of sum NDVI (RESTREND), 1981-2003




Figure 8. Confidence levels of RESTREND, 1981-2003




Figure 9. Global change in energy-use efficiency, 1981-2002




Figure 10. Global trend of positive climate-adjusted NDVI, 1981-2003




Figure 11. Confidence levels of positive climate-adjusted NDVI, 1981-2003

4 Land degradation and improvement

4.1 Land degradation

Land degradation means a loss of NPP but a decrease in NPP is not necessarily land degradation. To distinguish between declining productivity caused by land degradation and declining productivity due to other factors those other factors have to be reckoned with.

Rainfall variability has been accounted for using RUE-adjusted NDVI (Figure 5) and, also, by RESTREND (Figure 7).

We are unable to make allowance for changes in land use and management at the global level for lack of consistent time series data. In principle, land use change can be derived from changes in the shape of the annual NDVI wave; this is work in hand. Land use change will be addressed in forthcoming reports on particular *hot spots* and *bright spots*. See also Section 4.1.3.

The results are very different from the previous global assessment of land degradation (GLASOD) and challenge conventional wisdom. To answer the questions posed at the outset, comparisons were made with global data for land cover and land use, aridity, population density and, as proxies for poverty, infant mortality rates and proportion of underweight children under the age of five. The following discussion relates mainly to RUE-adjusted NDVI and its translation to NPP.

4.1.1 Which regions are hardest hit?

Areas severely affected (Table 1) include:

- Africa south of the Equator (13 per cent of global degrading area and 18 per cent of lost global NPP);
- Indo-China, Myanmar, Malaysia and Indonesia (6 per cent of the degrading area and 14 per cent of lost NPP);
- S China (5 per cent of the degrading area and 5 per cent of lost NPP);
- N-central Australia and parts of the western slopes of the Great Dividing Range (5 per cent of the degrading area and 4 per cent of lost NPP);
- The Pampas (3.5 per cent of the degrading area and 3 per cent of lost NPP);
- Swaths of the high-latitude forest belt in North America and Siberia.

The usual suspects – the dry lands around the Mediterranean, Middle East, South and Central Asia – are represented by only relatively small areas of degradation in southern Spain, the Maghreb, Nile delta, Iraqi marshes, and the Turgay steppe. Probably, many differences from the previous assessment arise because GLASOD compounded current land degradation with the legacy of centuries past. These are two different things; both are important; but most areas of historical land degradation have become stable landscapes – with a stubbornly low level of productivity. The present assessment deals only with 1981-2003 and we have no comparable data for earlier periods.

Table 1 presents country-by-country data for RUE-adjusted NDVI and NPP. The area data refer to pixels showing any declining trend - irrespective of degrees of confidence; by and large, the areas identified as high confidence are also those showing the most extreme trends - so intensity of degradation may be ranked more meaningfully according to total NPP loss than by gross degrading area.

Table 1: Statistics of degrading areas 1981-2003, by country
Countries or regions without degradation are not listed

Country	Degrading area (km ²)	% Territory	% global degrading area	Total NPP Loss (tonne C/23yr)	% total population	Affected people
Afghanistan	7658	1.17	0.025	62859	2.56	671770
Albania	2334	8.12	0.009	47250	4.29	137861
Algeria	63475	2.67	0.196	1977970	22.45	7168600
Andorra	281	60.00	0.001	2604	20.53	20865
Angola	828029	66.42	2.370	37602597	60.74	9263348
Argentina	902438	32.62	3.130	23556380	36.95	14455278
Armenia	743	2.49	0.003	13887	1.99	75632
Australia	1994268	25.94	6.182	46905279	11.31	2187493
Austria	28291	33.74	0.117	1835	21.51	1730745
Azerbaijan	2633	3.04	0.009	1230833	2.98	238076
Bahamas, The	4130	29.63	0.009	195146	32.01	19029
Bangladesh	68422	47.52	0.199	2851384	49.12	72728775
Belgium	5404	17.71	0.024	69560	13.48	1396093
Belize	3026	13.18	0.008	65978	16.94	39513
Benin	14155	12.57	0.041	373747	12.84	932170
Bhutan	27011	57.47	0.073	1705766	54.99	1332662
Bolivia	60339	5.49	0.175	1656319	16.39	1518038
Bosnia and Herzegovina	7737	15.13	0.030	157646	16.77	704321
Botswana	97831	16.30	0.284	4111881	30.74	476893
Brazil	1881702	22.11	5.381	63346318	26.67	46595573
Brunei	2663	46.15	0.008	127918	85.02	264401
Bulgaria	9139	8.24	0.035	178003	11.72	881122
Burkina Faso	9255	3.38	0.026	123795	8.26	1101414
Burundi	13516	48.56	0.037	972686	52.09	3881071
Belarus	4053	1.95	0.019	82416	2.56	254841
Cambodia	77958	43.06	0.225	2524942	24.03	3583464
Cameroon	151605	31.89	0.417	9657120	26.30	4326977
Canada	1985085	19.90	11.575	93963813	17.69	5509584
Cape Verde	375	9.30	0.001	12087	24.76	72997
Central African Republic	126927	20.37	0.356	3701988	23.27	894315
Chad	52735	4.11	0.152	627041	10.82	995721
Chile	77230	10.20	0.265	1950752	10.42	1645825
China	2193697	22.86	7.627	58840237	34.71	457202031
Colombia	291295	25.58	0.818	17999691	36.02	16309420
Comoros	181	8	0.001	17516	21.50	135144
Congo	201614	58.95	0.569	20091044	54.93	1895981
Costa Rica	14691	28.75	0.042	529400	13.41	592632
Croatia	2822	4.99	0.011	28610	7.95	338952
Cuba	32430	29.25	0.095	755492	28.31	3050838

Country	Degrading area (km ²)	% Territory	% global degrading area	Total NPP Loss (tonne C/23yr)	% total population	Affected people
Cyprus	266	2.87	0.001	9143	0.74	5164
Czech Republic	11218	14.22	0.048	304243	13.24	1358728
Denmark	91	0.21	0.001	290	0.24	10824
Djibouti	6107	27.76	0.017	19272	59.30	282700
Dominica	126	16.67	0.000	8976	7.57	4532
Dominican Republic	18507	37.98	0.054	560541	43.43	3843087
Ecuador	40136	14.15	0.101	2401058	16.13	2199904
Egypt	36514	3.65	0.112	16639	13.92	10100710
El Salvador	5585	26.54	0.016	234649	16.76	1139730
Equatorial Guinea	15376	54.81	0.037	1434524	45.39	171542
Eritrea	15573	12.84	0.045	33256	5.27	235381
Estonia	423	0.93	0.003	4083	0.75	9180
Ethiopia	296812	26.33	0.843	14276064	29.10	20650316
Falkland Islands (Islas Malvinas)	1635	13.43	0.009	50944	23.18	365
Finland	27779	8.24	0.178	327719	3.46	171458
France	46691	8.54	0.190	605160	10.48	6159286
French Guiana	24947	27.41	0.064	1033318	14.36	25745
Gabon	172865	64.58	0.471	23880	35.85	468972
Gambia, The	1396	12.35	0.004	26355	1.93	25821
Georgia	5647	8.10	0.021	141370	11.76	591918
Germany	32479	9.10	0.144	730980	6.97	5676882
Ghana	50365	21.11	0.143	2520819	20.95	4466773
Greece	6914	5.24	0.024	116915	6.76	662921
Guatemala	55884	51.32	0.163	2866596	30.46	3936416
Guinea	91415	37.18	0.262	2008342	46.51	4108349
Guinea-Bissau	18851	52.19	0.048	452425	43.43	536156
Guyana	93448	43.47	0.257	230119	26.49	198445
Haiti	11821	42.60	0.034	383261	34.56	2823765
Honduras	30145	26.89	0.084	1450818	23.38	1673952
Hungary	31398	33.75	0.128	765915	28.90	2810672
Iceland	34483	33.48	0.225	2693154	23.51	58021
India	592498	18.02	1.751	22484086	16.50	177437809
Indonesia	1028942	53.61	2.703	67679850	40.52	86656550
Iran	29190	1.77	0.095	282438	3.42	2572958
Iraq	28000	6.41	0.092	1030763	6.58	1718397
Ireland	6416	9.13	0.035	1363385	11.95	653134
Israel	3085	14.85	0.010	49570	30.07	2035012
Italy	28693	9.53	0.109	696409	7.80	4306062
Ivory Coast	117595	36.47	0.331	6221305	36.33	6252711
Jamaica	3372	30.68	0.010	106751	28.98	741313
Japan	130563	34.56	0.451	4268668	24.20	29666795
Jordan	13574	15.21	0.048	100582	19.13	1574810
Kazakhstan	487083	17.93	2.041	5308145	13.31	2131386
Kenya	104994	18.02	0.294	6612571	35.59	11803311
Korea, Peoples Republic of	60959	50.57	0.226	2206450	45.08	10124149
Korea, Republic of	54091	54.93	0.182	1570729	31.81	14364205
Kyrgyzstan	23189	11.68	0.087	282173	12.71	682075
Laos	133395	56.33	0.382	7232762	55.13	3304253

Country	Degrading area (km ²)	% Territory	% global degrading area	Total NPP Loss (tonne C/23yr)	% total population	Affected people
Latvia	4416	6.84	0.022	136363	9.49	213414
Lebanon	704	6.77	0.002	1894	3.37	123717
Lesotho	10344	34.08	0.033	485251	44.49	941131
Liberia	50500	45.34	0.123	2097992	38.12	1441085
Libya	12672	0.72	0.037	86083	6.92	402408
Lithuania	2664	4.09	0.016	55190	2.91	132351
Macedonia	1757	6.94	0.007	32910	1.42	30073
Madagascar	163843	27.91	0.492	6678189	21.56	3901784
Malawi	30869	26.05	0.089	1370895	19.89	2486085
Malaysia	175817	53.32	0.475	9257510	46.39	10401113
Mali	35637	2.87	0.106	357823	6.60	870031
Mauritania	6301	0.61	0.019	17918	2.18	67349
Mexico	487804	24.73	1.474	23871309	34.30	36234761
Moldova	1751	5.17	0.007	32362	3.17	133140
Mongolia	66559	4.25	0.271	623762	2.51	66138
Morocco	67399	15.09	0.201	2807952	35.71	11278600
Mozambique	226567	28.26	0.651	8398073	26.36	5155480
Myanmar (Burma)	358887	52.89	1.053	23625068	47.86	23608512
Namibia	288945	35.01	0.875	6388447	35.87	670983
Nepal	54704	38.85	0.182	2375267	48.93	13332932
Netherlands	7051	16.98	0.028	92199	17.25	2779551
New Caledonia	6902	36.21	0.020	1008271	31.44	48235
New Zealand	147014	54.72	0.545	6992963	30.97	1015925
Nicaragua	47223	36.47	0.134	2060424	29.28	1684227
Niger	22563	1.78	0.062	141699	6.61	844506
Nigeria	91443	9.90	0.256	3066735	13.33	17035650
Norway	57109	17.61	0.352	1212969	9.23	361786
Oman	419	0.20	0.002	3302	0.06	1848
Pakistan	20644	2.57	0.073	235711	3.58	5838072
Panama	8735	11.17	0.023	513509	7.78	232958
Papua New Guinea	205500	44.40	0.564	16275368	40.58	2019646
Paraguay	66704	16.40	0.200	1659008	66.97	4071629
Peru	197211	15.34	0.565	11414777	10.89	3001345
Philippines	132275	44.09	0.362	4100145	42.75	33064628
Poland	41514	13.28	0.188	890969	14.37	5505161
Portugal	11536	12.49	0.041	233458	4.58	440851
Puerto Rico	436	4.79	0.001	19231	2.91	111458
Reunion	175	6.98	0.001	6294	5.24	38724
Romania	16902	7.12	0.067	364407	4.47	980580
Russia	2802060	16.41	16.519	56663083	6.20	8588604
Rwanda	11404	43.30	0.031	1053147	39.11	3299059
Sao Tome and Principe	125	12.50	0.000	303560	21.82	28128
Saudi Arabia	8327	0.42	0.025	4335	2.00	471248
Senegal	34655	17.66	0.101	408832	20.49	2078643
Sierra Leone	35902	50.04	0.102	1507871	39.33	2103046
Singapore	243	37.50	0.001	5833	55.95	2017090
Slovakia	5066	10.37	0.021	110642	6.86	370606
Slovenia	2492	12.30	0.010	38132	17.99	396448
Solomon Islands	9065	31.86	0.030	628541	33.82	206290

Country	Degrading area (km ²)	% Territory	% global degrading area	Total NPP Loss (tonne C/23yr)	% total population	Affected people
Somalia	52520	8.24	0.149	1834048	14.77	1544921
South Africa	351555	28.82	1.124	23123364	38.14	17041101
Spain	63266	12.53	0.231	1712506	6.41	2417996
Sri Lanka	21057	32.09	0.060	634813	25.62	4788637
Sudan	166031	6.63	0.480	3627514	9.43	3280414
Suriname	50503	30.93	0.125	2102420	10.13	38529
Swaziland	16533	95.22	0.051	1226857	98.77	947510
Sweden	78964	17.55	0.475	1594303	10.37	841284
Switzerland	4982	12.07	0.020	106619	6.81	484619
Syria	11327	6.12	0.039	224233	6.71	1243265
Tajikistan	8412	5.88	0.030	104021	2.39	151676
Tanzania, United Republic of	386256	40.87	1.081	22603896	39.48	15300003
Thailand	309245	60.16	0.895	15990860	56.66	36991080
Togo	11064	19.48	0.032	2992723	12.79	654476
Trinidad and Tobago	675	13.16	0.002	113407	5.51	65120
Tunisia	12476	7.63	0.040	398423	15.47	1512817
Turkey	30851	3.95	0.111	453231	5.08	3571290
Turkmenistan	1273	0.26	0.005	8417	0.33	17554
Turks and Caicos Islands	92	21.43	0.001	15961	21.49	166
Uganda	41506	17.58	0.120	1513212	15.04	4112702
Ukraine	47414	7.85	0.200	1048460	5.25	2466172
United Kingdom	23506	9.60	0.103	262090	5.95	3324064
United States	1983886	20.60	7.935	39672698	10.79	31144568
Uruguay	87566	49.69	0.294	1874537	33.03	1058877
Uzbekistan	5974	1.34	0.022	123701	2.22	585887
Vanuatu	2210	14.97	0.005	4589	9.61	16965
Venezuela	207916	22.80	0.587	520023	8.28	2156456
Vietnam	134026	40.67	0.387	342632	35.27	28085074
Yemen	14422	2.73	0.032	7570	2.30	507751
Yugoslavia(Mace donia, Serbia, Montenegro)	10507	8.23	0.032	27197	6.37	678700
Zaire (Dem. Republic Congo)	1346914	57.43	3.760	3403930	53.49	32081359
Zambia	454630	60.41	1.312	19900481	50.07	5789865
Zimbabwe	180125	46.12	0.531	8861748	39.51	5424488
The World (land, excluding inland water body)	35058104	23.54	100.000	955221418	23.89	1537679148

The country rankings of severity of land degradation are: by percentage of the global degrading area, 1 Russia (16.5), 2 Canada (11.6), 3 USA (7.9) 4. China (7.6), and 5 Australia (6.2); by loss of NPP (million tonneC), 1 Canada (94), 2 Indonesia (68), 3 Brazil (63), 4 China (59), and 5 Australia (50); by percentage of the country affected, 1 Swaziland (95), 2 Angola (66), 3 Gabon (64), 4 Thailand (60), and 5 Zambia (60); and by rural population affected (millions), 1 China (457), 2 India (177), 3 Indonesia (86), 4 Bangladesh (72), and 5 Brazil (46). Table 2 shows the rankings of the LADA partner countries; China, Argentina and South Africa rank amongst the 20 most severely affected in terms of percentage area, loss of NPP and affected rural population. Each partner country is analysed individually in country reports (Bai and Dent 2007 a-f).

Table 2. Land degradation in LADA partner countries by global rank order

	% global area	NPP loss, million tonnesC	% country affected	affected rural population, million
China	4 (7.6)	4 (58.8)	(23)	1 (457)
Argentina	8 (3.1)	10 (23.6)	(33)	17 (14)
South Africa	15 (1.1)	11 (23.1)	(29)	14 (17)
Cuba	(0.09)	(0.8)	(29)	(3)
Senegal	(0.1)	(0.4)	(18)	(2)
Tunisia	(0.04)	(0.4)	(8)	(1.5)

4.1.2 *Is land degradation a global issue?*

Over the last 25 years, 24 per cent of the land area has been degrading (Table 1); this is on top of the legacy of thousands of years of mismanagement in some long-settled areas. GLASOD estimated that 15 per cent of the land was degraded (Appendix Table S1), and those areas are, by and large, not the same as the areas highlighted by the new analysis; land degradation is cumulative - this is the global issue.

Degrading areas currently support 1.5 billion rural people. In terms of C fixation, these areas represent a loss of NPP of 9.56×10^8 tonneC relative to the 1981-2003 mean; that is 9.56×10^8 tonneC *not* removed from the atmosphere - equivalent to 20 per cent of the global CO₂ emissions for 1980. At the shadow price for carbon used by the British Treasury in February 2008 (\$50/tonneC, Montbiot 2008) this amounts to \$US 48 billion in terms of lost C fixation. But the cost of land degradation is at least an order of magnitude greater in terms of C *emissions* from loss of soil organic carbon: as much as one third of the human-induced increase in atmospheric CO₂ and 20 per cent of global carbon emissions over the period 1989-1998 is related to land use change (IPCC 2000, Houghton 2008).

4.1.3 Is land degradation mainly associated with farming?

The short answer is *no*. Comparison of degrading areas with global land cover (JRC 2003) reveals that 18 per cent of degrading land is cropland (Table 3, codes 16-18), 23 per cent is broadleaved forest (codes 1-3), and 19 per cent needle-leaved forest (codes 4-5). Degradation is over-represented in both cropland and forest: cropland occupies only 12 per cent of the land area (and part of a further 4 per cent of mixed cover), broadleaved and needle-leaved forest together occupy 28 per cent of the land but 43 per cent of degrading land.

Comparison of degrading areas with global land use systems (FAO 2008, Tables 4 and 5) indicates that 47 per cent of degrading land is forest. Counter-intuitively, the proportion of degradation in the various forest categories is very similar: declining NPP is seen across 30 per cent of natural forest and supposedly protected forest, across 25-33 per cent of grazed forests and 33 per cent of plantations. At this stage, we can only speculate about the reasons: apart from land degradation as it is commonly understood, high-latitude *taiga* is subject to catastrophic fires and pest outbreaks that affect huge areas and, since the rate of recovery is slow, the 23-year period may encompass a whole cycle; and, surely, some of the recorded degradation is, in fact clearance for cropland and grazing. Twenty five per cent of the degradation is in grasslands (herbaceous vegetation in the FAO legend), where the natural and protected areas seem to be faring better than grazed areas.

Table 3. Global degrading and improving areas by land cover

<i>Code</i>	<i>Land cover</i>	<i>Total pixels (TP)</i> (0.54'x0.54')	<i>Degrading pixels (DP)</i> (0.54'x 0.54')	<i>DP/TP</i> (%)	<i>DP/TDP*</i> (%)	<i>Improving pixels (IP)</i> (0.54'x 0.54')	<i>IP/TP</i> (%)	<i>IP/TIP**</i> (%)
1	Tree Cover, broadleaved, evergreen	12875179	4222561	32.8	11.62	1985215	15.4	8.21
2	Tree Cover, broadleaved, deciduous, closed	8688097	2441119	28.1	6.72	877346	10.1	3.63
3	Tree Cover, broadleaved, deciduous, open	4099003	1616582	39.4	4.45	584110	14.3	2.41
4	Tree Cover, needle-leaved, evergreen	15080165	4633961	30.7	12.75	1020344	6.8	4.22
5	Tree Cover, needle-leaved, deciduous	8054159	2043323	25.4	5.62	427842	5.3	1.77
6	Tree Cover, mixed leaf type	5606446	993934	17.7	2.73	540412	9.6	2.23
7	Tree Cover, regularly flooded, fresh water	579763	228306	39.4	0.63	88405	15.2	0.37
8	Tree Cover, regularly flooded, saline water	115705	26157	22.6	0.07	17109	14.8	0.07
9	Mosaic: Tree Cover / Other natural vegetation	4269938	1097533	25.7	3.02	341516	8.0	1.41
10	Tree Cover, burnt	587270	225758	38.4	0.62	26659	4.5	0.11
11	Shrub Cover, closed-open, evergreen	3195387	1093184	34.2	3.01	226048	7.1	0.93
12	Shrub Cover, closed-open, deciduous	15605651	2953414	18.9	8.13	3263251	20.9	13.49
13	Herbaceous Cover, closed-open	17560702	2824775	16.1	7.77	3432708	19.5	14.19
14	Sparse herbaceous or sparse shrub cover	23573022	2567417	10.9	7.06	3115678	13.2	12.88
15	Regularly flooded shrub and/or herbaceous cover	3089962	689713	22.3	1.90	309398	10.0	1.28
16	Cultivated and managed areas	21692769	4522988	20.9	12.45	4306250	19.9	17.80
17	Mosaic: cropland/tree cover/other natural vegetation	4025653	1293550	32.1	3.56	559244	13.9	2.31
18	Mosaic: cropland/shrub and/or grass cover	3921904	692613	17.7	1.91	860554	21.9	3.56
19	Bare areas	24629888	931207	3.8	2.56	1641881	6.7	6.79
20	Water bodies	47103415	1105185	2.3	3.04	436082	0.9	1.80
21	Snow and ice	10660085	104790	1.0	0.29	99712	0.9	0.41
22	Artificial surfaces and associated areas	378999	35442	9.4	0.10	27944	7.4	0.12
23	No data	29056	120	0.4	0.0003	50	0.2	0.0002
Total		235422218	36343632		100	24187758		100.00

* TDP - total degrading pixels; **TIP – total improving pixels.

Table 4. Global degrading and improving areas by land use systems

<i>Code</i>	<i>Land use systems</i>	<i>Total pixels (TP)</i>	<i>Degrading pixels (DP)</i>	<i>DP/TP</i>	<i>DP/TDP¹</i>	<i>Improving pixels (IP)</i>	<i>IP/TP</i>	<i>IP/TIP²</i>
		(5'x5')	(5'x 5')	(%)	(%)	(5'x 5')	(%)	(%)
0	Undefined	272	12	4.4	0.003	11	4.0	0.004
1	Forestry - not managed (natural)	387251	115888	29.9	27.8	31553	8.1	11.4
2	Forestry - protected areas	78757	23425	29.7	5.6	8865	11.3	3.2
4	Forestry with pastoralism (moderate or higher)	128564	32872	25.6	7.9	15406	12.0	5.5
5	Forestry with pastoralism (moderate or higher), scattered plantations	24603	8149	33.1	2.0	3082	12.5	1.1
6	Forestry - scattered plantations	42757	13987	32.7	3.4	6301	14.7	2.3
7	Herbaceous - not managed /natural	212858	19012	8.9	4.6	17870	8.4	6.4
8	Herbaceous - protected areas	76707	10435	13.6	2.5	8560	11.2	3.1
9	Herbaceous with extensive pastoralism	154528	33253	21.5	8.0	31181	20.2	11.2
10	Herbaceous with moderate/ intensive pastoralism	129382	23199	17.9	5.6	33376	25.8	12.0
11	Herbaceous with intensive pastoralism	93193	19481	20.9	4.7	20471	22.0	7.4
13	Rain-fed agriculture (subsistence / commercial)	67231	17099	25.4	4.1	11060	16.5	4.0
14	Agro-pastoralism (moderate/ intensive)	81508	17232	21.1	4.1	17006	20.9	6.1
15	Agro-pastoralism (intensive)	126938	27282	21.5	6.6	25413	20.0	9.2
16	Agro-pastoralism (intensive with large-scale irrigation)	35170	6634	18.9	1.6	9235	26.3	3.3
17	Agriculture – large-scale irrigation (> 25% pixel size)	8464	2101	24.8	0.5	1491	17.6	0.5
18	Agriculture - protected areas	10551	2756	26.1	0.7	1704	16.2	0.6
19	Urban areas	52640	9114	17.3	2.2	6152	11.7	2.2

<i>Code</i>	<i>Land use systems</i>	<i>Total pixels (TP)</i> (5'x5')	<i>Degrading pixels (DP)</i> (5'x 5')	<i>DP/TP</i> (%)	<i>DP/TDP¹</i> (%)	<i>Improving pixels (IP)</i> (5'x 5')	<i>IP/TP</i> (%)	<i>IP/TIP²</i> (%)
20	Wetlands – not managed (natural)	35512	9060	25.5	2.2	3760	10.6	1.4
21	Wetlands - protected areas	5864	1336	22.8	0.3	782	13.3	0.3
22	Wetlands - mangroves	865	183	21.2	0.0	130	15.0	0.05
23	Wetlands - agro-pastoralism	331	58	17.5	0.0	87	26.3	0.03
24	Bare areas - not managed (natural)	225873	3737	1.7	0.9	5858	2.6	2.1
25	Bare areas - protected areas	74420	1332	1.8	0.3	1359	1.8	0.5
26	Bare areas - extensive pastoralism	67357	4719	7.0	1.1	7086	10.5	2.6
27	Bare areas – moderate or intensive pastoralism	32570	2012	6.2	0.5	5314	16.3	1.9
28	Water - Coastal or not managed (natural)	10283	603	5.9	0.1	476	4.6	0.2
29	Water - protected areas	9130	1468	16.1	0.4	650	7.1	0.2
30	Water - inland Fisheries	43480	9833	22.6	2.4	3445	7.9	1.2
100	Undefined	227	2	0.9	0.0005	0	0.0	0.0
Total		2217286	416274		100	277684		100

¹ TDP - total degrading pixels; ² TIP – total improving pixels.

Table 5. Global degrading/improving lands in the aggregated land use systems

<i>Land use system</i>	<i>Codes</i>	<i>Total pixels (TP)</i>	<i>Degrading pixels (DP)</i>	<i>DP/TP</i>	<i>DP/TDP¹</i>	<i>Improving pixels (IP)</i>	<i>IP/TP</i>	<i>IP/TIP²</i>
(LUS)		(5'x5')	(5'x 5')	(%)	(%)	(5'x 5')	(%)	(%)
Forestry	1-6	661932	194321	29.3	46.7	65207	9.9	23.5
Grassland	7-11	666668	105380	15.8	25.3	111458	16.7	40.1
Agricultural land	13-18	329862	73104	22.2	17.6	65909	20.0	23.7
Urban	19	52640	9114	17.3	2.2	6152	11.7	2.2
Wetlands	20-23	42572	10637	25.0	2.6	4759	11.2	1.7
Bare areas	24-27	400220	11800	2.95	2.8	19617	4.9	7.1
Water	28-30	62893	11904	18.9	2.9	4571	7.3	1.6
Undefined	0,100	499	14	2.8	0.0034	11	2.2	0.004

¹ TDP - total degrading pixels, ² TIP – total improving pixels

Degrading farmland makes up 18 per cent of the total degrading area shown in both the *LC2000* land cover map and the FAO *Land Use Systems* map. Interestingly, irrigated areas fare no better than the norm.

In Kenya over the period 1981-2003, NPP increased in woodland and grassland, but hardly at all in cropland; across 40 per cent of cropland it decreased - a critical situation in context of a doubling of human population over the same period (Bai and Dent 2006). In South Africa, NPP decreased overall and 29 per cent of the country suffered land degradation, including 41 per cent of all cropland (Bai & Dent 2007a); about 17 million people, 38 per cent of the South African population, depend on these degrading areas (Figure 12).




Figure 12. South Africa, land degradation and population affected, 1981-2003

Change of land use and management may generate false alarms about land degradation. Conversion of forest or grassland to arable, pasture or even perennial crops will usually result in an immediate reduction in NPP (and NDVI) but may well be profitable and sustainable, depending on management. Lack of consistent time series data for land use and management precludes a generalised analysis of land use change. This can be undertaken manually for the potential *hot spots* of land degradation, e.g. Chen and Rao 2008, and work is in hand on an automated procedure to map land use change from GIMMS NDVI data.

4.1.4 Is land degradation a dryland issue?

Drylands do not figure strongly in ongoing land degradation, apart from in Australia. Indeed, the recovery of the Sahel from the droughts of the 1980s is notable (Figure 2 and Olsson and others 2005). Globally, there is little between land degradation and Turc's aridity index correlation ($r = -0.12$); 78 per cent of degradation by area is in humid regions, 8 per cent in the dry sub-humid, 9 per cent in the semi-arid, and 5 per cent in arid and hyper-arid regions.

Similarly, in the three countries analysed (Cuba, Kenya, South Africa), there is no obvious correlation between degradation and soil and terrain. Indeed degradation is mostly on flat or gently-sloping areas; clearly this is a management issue.

4.1.5 Is it related to population pressure?

Comparison of rural population density (CEISIN 2007) with land degradation shows no simple pattern. Globally, the correlation coefficient is -0.3 ; in general, the more people the less degradation but, in some contexts, population pressure is positively related to land degradation. For South Africa (Figure 12, Figure 13), the correlation between land degradation and \log_e population density is positive ($r = 0.25$) but the former apartheid homelands have more than their fair share of degrading land (Bai and Dent 2007a) so something more than simple rural population density is at work.




Figure 13. South Africa: relationship between population density and land degradation / improvement

4.1.6 Is land degradation related to poverty?

Taking infant mortality rate and the percentage of children under five who are underweight (CEISIN 2007) as proxies, there is some global relationship between land degradation and poverty: correlation coefficients are 0.20 for both infant mortality and for underweight children. However, a much more rigorous analysis is

needed, especially to tease out the underlying biophysical and social and economic variables. This might be done using more specific geo-located data.

4.2 Land improvement

Land improvement is identified by: 1) a positive trend in RUE-adjusted sum NDVI and 2) a positive trend in energy-use efficiency, shown in Figure 10 as climate-adjusted NDVI. These areas account for 15.7 per cent of the land area; 18 per cent is cropland (20 per cent of the total croplands), 23 per cent is forest, and 43 per cent rangeland. Many gains in cropland are associated with irrigation but there are also swaths of improvement in rain-fed cropland and pastures in the Great Plains of North America, and in western India. Some of the NDVI gains are a result of increasing tree cover, either through forest plantations, especially in Europe and North America (FAO 2006), and some significant land reclamation projects, for instance in North China. However, some of the biomass gains may represent woodland and bush encroachment into rangeland and farmland - which is not generally regarded as land improvement.

We may attribute the general global increase in greenness to the increasing trends of atmospheric CO₂ concentrations and nitrate deposition. The increasing trend production across the Sahel probably includes an element of recovery from the devastating drought of the early 1980s, in spite of the adjustment for RUE and application of RESTREND. Increases in biomass in the Amazon basin may be related to lower rainfall, accompanied by decrease in growth-limiting cloudiness, but global data for net incoming radiation are not available to check this.

4.3 Urban areas

Whether urbanisation is degradation is arguable. Urbanisation brings a huge increase in the financial value of the land but, if it which involves sealing of the land surface, it is degradation according to our criterion of partial loss of loss of ecosystem function.

The CIESIN Global Rural Urban Mapping Project shows 3 per cent of the land area as urban. These areas are masked in the global maps. This makes only a small difference to the results: masking results in reduction of 0.5 per cent for the identified degrading land, and a reduction of 0.2 per cent for the improving land.

4.4 Comparison of indicators

Annual sum NDVI, representing annually accumulated greenness is our standard indicator of land degradation and improvement. Rain-use efficiency, RUE-adjusted NDVI and RESTREND are different ways of eliminating false alarms caused by rainfall variability (cf Sections 3.2.1 and 3.2.2, respectively). Each of the various measures is useful in its own right. Both NDVI (as an index of greenness) and rain-

use efficiency (production per unit of rainfall) are simple and well known concepts. An advantage of RUE-adjusted NDVI is that, for areas considered as degrading or improving, the original NDVI values are retained. These can then be translated into NPP, which is open to economic analysis - e.g. in terms of loss or gain in carbon fixation.

Comparing the various indicators: the global trends of sum NDVI and RESTREND are almost identical (Figure 2 and Table 6): about 54 per cent of land area shows positive trend in both sum NDVI and RESTREND, 26 per cent show negative change and 20 per cent, no change. If we take negative RUE-adjusted NDVI as the primary indicator of degradation, then 96 per cent of degrading land, so defined, also shows negative RESTREND. If we take a positive RUE-adjusted NDVI as the primary indicator of improvement, then nearly all of the improving land, so defined, also shows positive RESTREND.

Comparing RUE with RESTREND, 15 per cent of the land area shows negative trend in both RUE and RESTREND, 37 per cent shows positive trend in both RUE and RESTREND and 20 per cent no change. But we get mixed signals from the remaining 28 per cent: either positive RUE and negative RESTREND, or vice versa. Comparing RUE and RESTREND on the same basis as above, 69 per cent of the degrading land shows both negative RESTREND and negative RUE; 90 per cent of the improving land shows positive RESTREND and positive RUE.

Table 6. Global comparison of trends in various indicators

Indicators	Total pixel (%)	Negative trend (%)	Positive trend (%)	No change (%)	Mixed (%)
Annual sum NDVI	100	25.7	54.0	20.3	0
RESTREND ¹	100	25.8	53.2	21.0	0
RUE	100	31.1	48.1	20.8	0
Sum NDVI \cap RESTREND	100	23.7	51.3	20.9	4.1
Sum NDVI \cap RESTREND with LI ²			99.5		
Sum NDVI \cap RESTREND with LD ³		96.3			
RUE \cap RESTREND	100	15	36.9	20.3	27.8
RUE \cap RESTREND with LI			91.3		
RUE \cap RESTREND with LD		68.5			

¹ Residual trend of sum NDVI; ² LI - identified improving land; ³ LD - identified degrading land.

Overall, RUE-adjusted NDVI provides a more restricted delineation of areas of degradation, as may be seen from Figure 14 which compares the four indicators for part of South Africa.




Figure 14. Eastern South Africa: sum NDVI, RUE-adjusted NDVI, rain-use efficiency, and RESTREND

5 What GLADA can and cannot show; further analysis

We have defined land degradation as a long-term loss of ecosystem function and use net primary productivity (NPP) as an indicator. GLADA is an interpretation of GIMMS NDVI data – which is taken as a proxy for NPP. The proxy is several steps removed from the symptoms of land degradation as it is commonly understood – such as soil erosion, salinity, or nutrient depletion. The same goes for land improvement. Any particular trend in NDVI may mean something very different in Central Africa from a quantitatively similar trend in South China or South Africa – both in terms of the kind of land degradation and the underlying causes.

A declining trend of NPP, even allowing for climatic variability, may not even be reckoned as land degradation: urban development is generally considered to be development - although it brings a loss of ecosystem function; land use change from forest or grassland to cropland of lesser biological productivity may or may not be accompanied by soil erosion, compaction and nutrient depletion - and it may well be sustainable and profitable, depending on management. Similarly, an increasing trend of NPP means greater biological production but may reflect, for instance, bush encroachment in rangeland or cropland - which is not land improvement as commonly understood.

Land use change is clearly a major driver of change in NPP. However, we have not been able to make allowance for land use change in the same way as for rainfall and temperature because available data are not compatible with the GIMMS data. Analysis of land use change based on the GIMMS dataset is in hand but, in the interim, we may refer to the trends in annual minimum and standard deviation NDVI (Figures S1 and S6) as a indicators of the areas most affected by land use change.

The 8km resolution of the GIMMS data is a limitation in two senses. First, an 8km pixel integrates the signal from a wider surrounding area. Many symptoms of even very severe land degradation, such as gulleys, rarely extend over such a large area; they must be severe indeed to be seen against the signal of the surrounding unaffected areas. More detailed analysis is possible for those areas that have higher resolution time series data, notably South Africa (Wessels and others 2004). Secondly, an 8km pixel or even a 1km pixel cannot be checked by a windscreens survey; and a 23-year trend cannot be checked by a single snapshot.

As already mentioned, the lack of consistent time series data on land use prevents a general accounting of land use change in the global assessment.

Some inherent limitations of the NDVI data have already been flagged: saturation of the NDVI signal by dense vegetation - leading to a lack of precision for forest mapping; interference by cloud in perennially cloudy areas; and the scant rainfall observations in many parts of the world.

1. A new GIMMS dataset, further corrected and updated to 2006 is now available; an updated VASCLimO dataset is also expected later this year; this will enable updating of the present analysis.
2. As an indicator of land degradation and improvement, f_{PAR} is preferred to NDVI – in its own right as a direct measurement of an important biophysical parameter, and to derive NPP through either the MODIS or JRC model. Data are available from year 2000 and at 1km resolution rather than the 8km resolution of GIMMS. Looking forward, these data would be preferred for monitoring and early warning;
3. Rather than using sometimes-sparse station-observed data, rainfall modelled from earth-observation satellite data are now available at the same level of precision as f_{PAR} data, e.g. TRMM (2008) and WaterWatch (2008). Again, this is preferred for the future but not applicable to the present analysis;
4. Cloud interference may be minimised by calculating trends for longer time steps, up to five years rather than an annual. This entails loss of precision.

The present analysis used only a fraction of the information available in the GIMMS data:

1. We have used simple linear regression of the 23-year GIMMS period to analyse the trends of NDVI and NDVI derivatives. It is possible to use power functions and separate, say successive 10-year periods;
2. There is valuable information in the seasonal shape of the NDVI curves that may be analysed, e.g. by harmonic analyses of NDVI time series (HANTS, de Wit 2004); critical information on timing of changes in land use and management can be interpreted manually from time series for individual pixels but algorithm development is required for regional and global application;
3. Visualization can be greatly improved by three-dimensional overlays of the NDVI/NPP trend surfaces over topography and in combination with other data layers;
4. Comparison of the present situation with potential biological productivity *without* human-induced land use change – the Garden of Eden scenario – modelled from climatic, soils and topographic data using, e.g. the BIOME-BGC model (Thornton and others 2005), will enable separation of the last 25 years of land degradation from the historical legacy.

In the next phase of GLADA, individual areas of land degradation and improvement, identified on the basis of NDVI indicators, will be characterised by geo-location of GLADA indicators alongside DEM, land use and soils data. Within individual soil and terrain (SOTER) units, a more refined analysis will be performed by modelling the relationships between rainfall, temperature and NDVI for the whole SOTER unit, then calculating the deviation for individual pixels. This makes allowance for soil and terrain and lessens the bias of the rainfall-NDVI model imposed by that pixel's deviations from the general trend.

At the same time, the continuous field of the index of land degradation derived from NDVI and climatic data will enable a statistical examination of other data for which continuous spatial coverage is not available - for instance spot measurements of soil attributes, and other social and economic data that may

reflect the drivers of land degradation, provided that these other data are geo-located.

Time series of 30m-resolution Landsat imagery will be used to identify land use the probable kinds of land degradation. Finally, field examination of the identified areas of degradation and improvement will be undertaken by national teams within the wider LADA program. A rigorous procedure is laid out in the forthcoming *LADA Field Handbook*. Apart from systematically and consistently characterising the situation on the ground across a range of scales, the field teams should address the following questions:

1. Is the biomass trend indicated by GLADA real?
2. If so, does it correspond with physical manifestations of land degradation or improvement that are measurable on the ground?
3. In the case of a negative answer to either of the preceding questions, what has caused the observed NPP trend? For instance, is it a question of timing of observations - where the situation on the ground has recovered?

6 Conclusions

1. **Land degradation and improvement have been assessed by remotely sensed indicators of biomass productivity.** The indicators show clear regional trends over the period 1981-2003, both decreasing and increasing, which may be interpreted as land degradation or improvement, respectively.
2. **Biomass trends depend on several factors other than land degradation and improvement:**
 - a. **We have taken account of rainfall variability in two ways: by screening NDVI trends for rain-use efficiency (RUE) in those areas where productivity is limited by rainfall, and by residual trends analysis.** RUE (net primary productivity per unit of rainfall), accounts for rainfall variability and, to some extent, local soil and land characteristics. We assume that, where NPP is limited by rainfall, a declining trend in RUE indicates land degradation. Where rainfall is not limiting, NPP is the best indicator available. Taken together, the two indicators may provide a more robust assessment than either used alone. As well as RUE, we calculated residual trends of NDVI, which point in the same direction as RUE-adjusted NDVI;
 - b. **Energy-use efficiency** (the ratio of NPP and accumulated temperature) proves to be more of an issue in defining improving areas than degrading areas;
 - c. **Potentially significant factors for which there are no consistent global data include changes in land use and management** and net incoming radiation.
3. **All changes measured by climate-adjusted NDVI/NPP are not land degradation or improvement as usually understood.** Change of land use from forest to cropland of lesser biological productivity, or an increase in grazing pressure, or a market adjustment to a less-intensive management will all decrease NDVI. These changes may or may not be accompanied by soil erosion, salinity, or other symptoms of land degradation of concern to soil scientists. In the same way, pastoralists will not consider bush encroachment as land improvement although it may increase biomass; these are limitations of a proxy indicator. Also, ambiguous data from the boreal forest belt may reflect catastrophic forest fires and outbreaks of pests, for instance the mountain pine beetle (Kurz and others 2008). These are part of the natural cycle but ecosystem recovery is slow in cold and dry regions as well as in dry regions and events falling towards the end of the 23-year measurement period affects the measured trend. This may not be land degradation - we should expect recovery - but if these events are themselves be related to climate change the system may not recover.
4. **GLADA presents a different picture from previous assessments of land degradation which compounded historical land degradation with what is happening now.** The data since 1981 indicate current trends but tell us

nothing about the historical legacy. For many purposes, it is more important to address on-going land degradation; much historical land degradation may be irreversible.

5. **As a quantitative measure of land degradation, loss of NPP has been calculated for those areas where *both* NPP and RUE are declining.** This is likely to be a conservative estimate since globally, NPP has increased over the period. Also, where NPP is increasing but RUE is declining, some process of land degradation may have begun that is reducing NPP but is not yet reflected in declining NPP.
6. By the same reasoning, **RUE should be used alone for *early warning* of land degradation, or a herald of improvement.** Where NPP is rising but RUE declining, some process of land degradation might be under way that is not yet reflected in declining NPP; it will remain undetected if we consider only those areas where both indices are declining. The reverse also holds true: we might forgo promising interventions that increase RUE but have not yet brought about increasing NPP.
7. **Long-term trends of NDVI derivatives are unsophisticated indicators of land degradation and improvement. The various kinds of land degradation and improvement are not distinguished but as a proxy, NDVI/NPP trend does provide a globally consistent yardstick, and it does highlight places where biologically significant change is happening.** And this is its purpose: in the parent LADA program, this global scan will be used to direct attention to areas that demand investigation and action on the ground.

Acknowledgements

This work is part of the GEF/UNEP/FAO project *Land Degradation Assessment in Drylands*.

We thank CJ Tucker, JE Pinzon and ME Brown for access to the GIMMS dataset; J Grieser for the VASCLIM0 precipitation data, T Fuchs for the GPCC precipitation data, and M Salmon for the CRU climatic data; we are indebted to A Anyamba, R Biancalani, F Nachtergaele, SG Prince, BGJS Sonneveld, A Tengberg, GWJ van Lynden, P Vlek, and KJ Wessels for critical review; to P Tempel for help with data handling, G Heuvelink for help with statistics, and to YGL Karpes-Liem for text layout.

References

- Alexandrov GA & T Oikawa 1997 Contemporary variations in terrestrial net primary productivity: the use of satellite data in the light of the extremal principle. *Ecological Modelling* 95, 113-118
- Anyamba A & CJ Tucker 2005 Analysis of Sahelian vegetation dynamics using NOAA-AHRR NDVI data from 1981-2003. *Journal of Arid Environments* 63, 596-614
- Asrar GM, M Fuchs, ET Kanemasu & JL Hatfield 1984 Estimating absorbed photosynthetically active radiation and leaf area index from spectral reflectance in wheat. *Journal of Agronomy* 87, 300-306
- Bai ZG & Dent DL 2006 *Global assessment of land degradation and improvement: pilot study in Kenya*. Report 2006/01, ISRIC - World Soil Information, Wageningen
- Bai ZG & DL Dent 2007a *Land degradation and improvement in South Africa 1. Identification by remote sensing*. Report 2007 /03, ISRIC - World Soil Information, Wageningen
- Bai ZG & DL Dent 2007b *Land degradation and improvement in Cuba 1. Identification by remote sensing*. Report 2007/04, ISRIC - World Soil Information, Wageningen
- Bai ZG & DL Dent 2007c *Land degradation and improvement in China 1. Identification by remote sensing*. Report 2007/06, ISRIC - World Soil Information, Wageningen
- Bai ZG & DL Dent 2007d *Land degradation and improvement in Argentina 1. Identification by remote sensing*. Report 2007/05, ISRIC - World Soil Information, Wageningen
- Bai ZG & DL Dent 2007e *Land degradation and improvement in Senegal 1. Identification by remote sensing*. Report 2007/07, ISRIC - World Soil Information, Wageningen
- Bai ZG & DL Dent 2007f *Land degradation and improvement in Tunisia 1. Identification by remote sensing*. Report 2007/08, ISRIC - World Soil Information, Wageningen
- Bai ZG 2006 *Assessing land degradation in the Chittagong Hill Tracts, Bangladesh, using NASA GIMMS*. Report 2006/6, ISRIC - World Soil Information, Wageningen
- Bai ZG, Dent DL. & Schaepman ME 2005 *Quantitative global assessment of land degradation and improvement: pilot study in North China*. Report 2005/6, ISRIC - World Soil Information, Wageningen
- Beck C, J Grieser & B Rudolf 2005 A new monthly precipitation climatology for the global land areas for the period 1951-2000. 181-190 in *Climate Status Report 2004*. German Weather Service, Offenbach
- Brown ME, JE Pinzon & K Didan 2006 Evaluation of the consistency of long-term NDVI time series derived from AVHRR, SPOT-Vegetation, SeaWiFS, MODIS and Landsat ETM+. *IEEE Transactions Geoscience Remote Sensing* 44, 1787-1793

- CEISIN 2007 Socioeconomic Data Applications Center, Center for International Earth Science Information Network, Columbia University, Palisades NY <http://sedac.ciesin.columbia.edu/gpw>;
http://sedac.ciesin.columbia.edu/povmap/ds_global.html (Accessed October 10, 2007)
- Chen S & P Rao 2008 Land degradation monitoring using multi-temporal Landsat TM/ETM data in a transition zone between grassland and cropland of northeast China. *International Journal of Remote Sensing* doi: 10.1080/01431160701355280.
- CIESIN 2004 *Global Rural-urban mapping project: urban/rural extents*. Center for International Earth Science Information Network, Columbia Univ., Palisades NY
- Engelen VWP van & TT Wen 1995 *Global and national soils and terrain databases (SOTER). Procedures manual (revised edition)*. UNEP-FAO-ISRIC, Wageningen
- FAO 2006 *Global forest resources assessment 2005*. Forestry paper 147 FAO, Rome
- FAO 2008 Land use systems of the World. <http://www.fao.org/geonetwork/srv/en/metadata.show>. Accessed 15 June 2008
- Fensholt R, I Sandholt & MS Rasmussen 2004 Evaluation of MODIS LAI, fAPAR and the relation between fAPAR and NDVI in a semi-arid environment using in situ measurements. *Remote Sensing of Environment* 91, 490-507
- Fensholt R, I Sandholt, MS Rasmussen, S Stison & A Diouf 2006 Evaluation of satellite-based primary production modelling in the semi-arid Sahel. *Remote Sensing of the Environment* 105 173-188
- Gebremichael M & AP Barros 2006 Evaluation of MODIS gross primary productivity (GPP) in tropical monsoon regions. *Remote Sensing of Environment* 100, 150-166
- Hall F, J Masek & GJ Collatz 2006 Evaluation of ISLSCP Initiative II FASIR and GOMMS NDVI products and implications for carbon cycle science. *Journal of Geophysical Research* 111: D22S08 doi:10.1029/2006JD007438
- Holm AMS, W Cridland & ML Roderick 2003 The use of time-integrated NOAA NDVI data and rainfall to assess landscape degradation in the arid shrubland of Western Australia. *Remote Sensing of Environment* 85, 145-158
- Houérou HN le 1984 Rain-use efficiency: a unifying concept in arid-land ecology. *Journal of Arid Environments* 7, 213-247
- Houérou HN le, RL Bingham & W Skerbek 1988 Relationship between the variability of primary production and variability of annual precipitation in world arid lands. *Journal of Arid Environments* 15, 1-18
- Houghton RA 2008 Carbon flux to the atmosphere from land use changes 1850-2005. *TRENDS: A compendium of data on global change*. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Dept Energy, Oak Ridge TE <http://cdiac.ornl.gov/trends/landuse/hughton.html> Accessed 13 May 2008
- Illius AW and TG O'Connor 1999 On the relevance of nonequilibrium concepts to arid and semiarid grazing systems. *Ecological Applications* 9, 798-813
- IPCC 2000 Global perspective. 25-51 In: Watson RT, IR Noble, B Bolin and others (editors) *Land use, land-use change, and forestry*. Published for the Intergovernmental Panel on Climate Change by Cambridge University Press, Cambridge

- ISRIC 2008 a *Global SOTER landform classification scale 1:1 million*. ISRIC Working Paper, Wageningen
- ISRIC 2008b *Enhanced soil information for LADA partner countries, scale 1:1million*. ISRIC Working Paper, Wageningen
- Jones JAA 1997 *Global hydrology: processes, resources and environmental management*. Longman, Harlow
- JRC 2003 *Global Land Cover 2000 database*. European Commission, Joint Research Centre, <http://www-gem.jrc.it/glc2000>. Accessed February 2, 2007
- Justice CO, JRG Townshend, EF Vermote and others 2002 An overview of MODIS Land data processing and product status. *Remote Sensing of Environment* 83, 3-15
- Justice CO, G Dugdale, JRG Townshend and others 1991 Synergism between NOAAVHRR and Meteosat data for studying vegetation development in semi-arid West Africa. *International Journal of Remote Sensing* 12, 1349-1368
- Kurz WA, CC Dymond, G Stinson and others 2008 Mountain pine beetle and forest carbon feedback to climate change. *Nature* 452, 24, 987 doi: 10.1038/nature06777
- Livezy RE & WY Chen 1983 Statistical field significance and its determination by Monte Carlo techniques. *Monthly Weather Review* 111, 46-59
- Mitchell TD & PD Jones 2005 An improved method of constructing a database of monthly climate observations and associated high-resolution grids. *International Journal of Climate* 25, 693-712
- Monbiot G 2008 It makes economic sense to kill people. *The Guardian Weekly* 20.02.08, page 19
- Myneni RB, R Ramakrishna, R Nemani & S Running 1997 Estimation of global leaf area index and absorbed PAR using radiative transfer models. *IEEE Transactions Geoscience Remote Sensing* 35 1380-1393
- Nemani RR, CD Keeling, H Hashimoto and others 2003 Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science* 300, 1560-1563
- O'Connor TG, LM Haines & HA Snyman 2001 Influence of precipitation and species composition on phytomass of a semi-arid African grassland. *Journal of Ecology* 89, 850-860
- Oldeman LR, RTA Hakkeling & WG Sombroek 1991. *World map of the status of human-induced soil degradation, 2nd edition*. ISRIC, Wageningen
- Olsson L, L Eklundh & J Ardö 2005. A recent greening of the Sahel – trends, patterns and potential causes. *Journal of Arid Environments* 63, 556-566
- Plummer S 2006 On validation of the MODIIS gross primary production product. *IEEE Transactions on Geoscience and Remote Sensing* 44, 7, 1936-1938
- Prince SD, KJ Wessels, CJ Tucker & SE Nicholson 2007 Desertification in the Sahel: a reinterpretation of a reinterpretation. *Global Change Biology* 13, 1308-1313
- Rasmussen MS 1998a Developing simple, operational, consistent NDVI-vegetation models by applying environmental and climatic information Part II: Crop yield assessment. *International Journal of Remote Sensing* 19, 119-139
- Rasmussen MS 1998b Developing simple, operational, consistent NDVI-vegetation models by applying environmental and climatic information: Part I: Assessment of net primary production. *International Journal of Remote Sensing* 19, 97-117

- Ripple WJ 1985 Asymptotic reflectance characteristics of grass vegetation. *Photogrammetric Engineering Remote Sensing* 51, 1915-1921
- Running SW, FA Heinsch, M Zhao, M Reeves & H Hashimoto 2004 A continuous satellite-derived measure of global terrestrial production. *Bioscience* 54, 547-560
- Seaquist JW, L Olsson & J Ardö 2003 A remote sensing-based primary production model from grassland biomes. *Ecological Modelling* 169, 131-155
- Schneider U, T Fuchs, A Meyer-Christoffer & B Rudolf 2008 *Global precipitation analysis products of the GPCC, full data reanalysis product version 4*. Global Precipitation Climatology Centre, Deutscher Wetterdienst, Offenbach ftp://ftp-anon.dwd.de/pub/data/gpcc/html/fulldata_download.html accessed August 8 2008
- Snyman HA 1998 Dynamics and sustainable utilization of rangeland ecosystems in arid and semi-arid climates of southern Africa. *Journal of Arid Environments* 39, 645-666
- Sonneveld BGJ. & DL Dent 2007 How good is GLASOD? *Journal of Environmental Management* doi:10.1016/j.jenvman.2007.09.008
- Thornton PE, Running SW & Hunt ER 2005 *Biome-BGC: Terrestrial Ecosystem Process Model, Version 4.1.1*. Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge TE <http://www.daac.ornl.gov>
- TRMM 2008. <http://trmm.gsfc.nasa.gov/> accessed May 1, 2008
- Tucker CJ, JE Pinzon & ME Brown 2004 *Global Inventory Modeling and Mapping Studies (GIMMS) satellite drift corrected and NOAA-16 incorporated normalised difference vegetation index (NDVI), monthly 1981-2002*. University of Maryland
- Turner DP, WD Ritts, WB Cohen and others 2003 Scaling gross primary production (GPP) over boreal and deciduous forest landscapes in support of MODIS GPP product validation. *Remote Sensing of Environment* 88, 256-270
- Turner DP, WD Ritts, WB Cohen and others 2006 Evaluation of MODIS NPP and GPP products across multiple biomes. *Remote Sensing of Environment* 102, 282-292
- UNCED 1992 Managing fragile ecosystems: combating desertification and drought Agenda 21, Chapter 12 United Nations Conference on Environment and Development <http://www.un.org/esa/sustdev/documents/agenda21>. Accessed August 5, 2005
- UNEP 2007 *Global Environmental Outlook GEO-4*, UN Environment Programme, Nairobi
- WaterWatch 2008 <http://www.WaterWatch.nl> accessed May 1, 2008
- Wessels KJ, SD Prince, J Malherbe and others 2007 Can human-induced land degradation be distinguished from the effects of rainfall variability? A case study in South Africa. *Journal of Arid Environments* 68, 271-297
- Wessels KJ, SD Prince, PE Frost & D VanZyl 2004 Assessing the effects of human-induced land degradation in the former homelands of northern South Africa with a 1-km AVHRR NDVI time-series. *Remote Sensing and Environment* 91, 47-67
- Wit de A 2004 The IDL-ENVI implementation of the HANTS algorithm (personal communication)

Zhao M, FA Heinsch, RR Nemani & SW Running 2005 Improvements of the MODIS terrestrial gross and net primary production global data set. *Remote Sensing of Environment* 95, 164-176

Zhao M, SW Running & RR Nemani 2006 Sensitivity of Moderate Resolution Imaging Spectroradiometer (MODIS) terrestrial primary production to the accuracy of meteorological reanalyses. *Journal of Geophysical Research* 111, G01002, doi:10.1029/2004JG000004

Appendix 1: Data

Table S1 GLASOD estimates of human-induced soil degradation, million ha

Kind of degradation	World	Asia	West Asia	Africa	Latin America and Caribbean	North America	Australia and Pacific	Europe
Water erosion	1094	440	84	227	169	60	83	115
Wind erosion	548	222	145	187	47	35	16	42
Nutrient depletion	135	15	6	45	72	-	+	3
Salinity	76	53	47	15	4	-	1	4
Contamination	22	2	+	+	+	-	-	19
Physical	79	12	4	18	13	1	2	36
Other	10	3	1	2	1	-	1	2
Sum	1964	747	287	494	306	96	103	218

GLASOD, reporting in 1991, indicated that 15 per cent of land was degraded. The highest proportions were reported for Europe (25 per cent), Asia (18 per cent) and Africa (16 per cent); the least in North America (5 per cent). By the same measure, as a proportion of the degraded area, soil erosion affected 83 per cent of the global degraded area (ranging from 99 per cent in North America to 61 per cent in Europe); nutrient depletion affected 4 per cent globally but 28 per cent in South America; salinity less than 4 per cent worldwide but 16 per cent in West Asia; chemical contamination about 1 per cent globally but 8 per cent in Europe; soil physical problems 4 per cent globally but 16 per cent in Europe.

Table S2 Statistics of NDVI indicators*

NDVI indicators	NDVI values			Pixels (%)		% NDVI change/year			Δ NDVI/year		
	min	max	mean	Pos.	Neg.	Pos.	Neg.	mean	Pos.	Neg.	mean
Minimum	0.132	0.296	0.182	59.5	40.5	0.966	0.851	0.232	0.0019	0.0014	0.00055
Maximum	0.527	0.716	0.622	50.5	49.5	0.391	0.260	0.060	0.0019	0.0017	0.00014
Max-Min	0.322	0.575	0.443	46.0	54.0	0.732	0.670	-0.036	0.0023	0.0026	-0.00039
Mean	0.314	0.411	0.365	67.6	32.4	0.368	0.245	0.164	0.0011	0.0009	0.00043
Sum	3.762	4.932	4.383	67.6	32.4	0.368	0.245	0.164	0.0129	0.0104	0.00518
STD	0.110	0.192	0.150	53.2	46.8	0.707	0.684	0.046	0.0008	0.0008	0.00005
CoV	0.206	0.394	0.294	41.5	58.5	0.772	0.735	-0.113	0.0036	0.0033	-0.00044

*In the calculations of the min., max. and mean values of each NDVI indicator, an average value of the all pixels in the vegetated area, defined as areas with net primary productivity greater than 1 g C m⁻² year⁻¹, were calculated. For example, min. value of the Maximum NDVI indicator: overlay statistic minimum of CELL STATISTIC in ArcMap was performed to extract minimum values of the time series annual Maximum NDVI for each pixel over the period (1981-2003), and the averaged minimum value of the maximum NDVI for all pixels was assigned as min. for the Maximum NDVI indicator; max. value of the Maximum NDVI indicator: overlay statistic maximum of CELL STATISTIC in ArcMap was performed to extract maximum values of the time series annual Maximum NDVI for each pixel over the period (1981-2003), and the averaged maximum value of the maximum NDVI for all pixels was assigned as max. for the Maximum NDVI indicator; mean value of the Maximum NDVI indicator: overlay statistic mean of CELL STATISTIC in ArcMap was performed to extract mean values of the time series annual Maximum NDVI for each pixel over the period (1981-2003), and the averaged mean value of the maximum NDVI for all pixels was assigned as mean for the Maximum NDVI indicator.

The rates of the positive and negative pixels were counted from the slope of the regression, i.e., positive slope (pos.) negative slope (neg.).

% NDVI change/year was calculated from the trend maps for each NDVI indicator: positive value (pos.) is the average of the all pixels with a positive trend; negative (neg.) is the average of the all pixels with a negative trend; mean value is the average of the all pixels; Δ NDVI/year is calculated the same as % NDVI change but from the absolute change maps.

Appendix 2: Analytical methods

Derivation of NDVI indicators

ArcGIS Spatial Analyst, ERDAS IMAGINE and ENVI-IDL were used to calculate NDVI minimum, maximum, maximum-minimum, mean, sum, standard deviation, and coefficient of variation (Appendix 3), as well as climate variables. The fortnightly NDVI data were geo-referenced and averaged to monthly; annual NDVI indicators (for the calendar year in the northern hemisphere and from 1 October to following 30 September for the southern hemisphere) were derived for each pixel; their temporal trends were determined by linear regression at an annual interval and mapped to depict spatial changes.

A negative slope of linear regression indicates a decline of green biomass and a positive slope, an increase – except for STD and CoV which indicate trends in variability. The absolute change (Δ in map legends, titled “changes in”) is the slope of the regression; the relative change (% in map legends, titled “trend in”) is $100(\text{slope of the regression}/\text{multi-year mean})$.

Monthly grids of rainfall for the period 1981-2002 were geo-referenced and re-sampled to the same spatial resolution as the NDVI (8km) using neighbourhood statistics. Spatial pattern and temporal trend of rainfall and rain-use efficiency (RUE, the ratio of annual NDVI and annual rainfall) for each pixel were determined by regression.

Land degradation was identified by negative trends of both biomass and rain-use efficiency. To distinguish between declining productivity caused by land degradation, and declining productivity due to other factors, rainfall variability has been accounted for by, first, identifying pixels where there is a positive relationship between productivity and rainfall. Secondly, for those areas where productivity depends on rainfall, rain-use efficiency has been considered: where productivity declined but RUE increased, we attribute the decline of productivity to declining rainfall and those areas are masked.

Land improvement was identified by positive changes in sum NDVI, positive rain-use efficiency in those areas where there is which has a positive correlation between sum NDVI and rainfall and RUE, and positive energy-use efficiency.

Plots of both land degradation and land improvement were masked by the mapped urban extents

Statistical tests

The trend analysis assumes that the data are spatially and temporally independent. This was tested by examining autocorrelation coefficients following Livezy and Chen (1983). When the absolute values of the autocorrelation coefficients of lag-1 to lag-3 calculated for a time series consisting of n observations are not larger than the typical critical value corresponding to 5 per cent significance level, i.e., $1.96/\sqrt{n}$, the observations in this time series can be accepted as being independent from each other.

The T-test was used to arrange the slope values in classes showing strong or weak positive or negative trends:

$$T = b / se(b)$$

Where b is the calculated slope of the regression line between the observation values and time and $se(b)$ represents the standard error of b .

The class boundaries were defined for 95 per cent confidence level; trends were labelled *high* if the T -values of the slope exceeded the 0.025 p -value of either tail of the distribution; lesser T - values were labelled *low*.

In addition, SPSS and MS Excel were employed to analyze trends, correlations and significances of the non-gridded variables.

Maps of the degrading areas or improving areas were overlaid on the other maps. Corresponding comparative values were calculated, pixel-by-pixel and a univariate correlation calculated.

Appendix 3: NDVI indicators of the land degradation / improvement

Minimum NDVI: The lowest value that occurs in any one year (annual) - which is usually at the end of the dry season. Variation in minimum NDVI may serve as a baseline for other parameters.

Maximum or peak NDVI: Represents the maximum green biomass. The large spatial variations reflect the diverse landscapes and climate.

Maximum-minimum NDVI: The difference between annual maximum and minimum NDVI reflects annual biomass productivity for areas with one, well-defined growing season but may not be meaningful for areas with bimodal rainfall.

Sum NDVI: The sum of fortnightly NDVI values for the year most nearly aggregates annual biomass productivity.

Standard deviation (STD): NDVI standard deviation is the root mean square deviation of the NDVI time series values (annual) from their arithmetic mean. It is a measure of statistical dispersion, measuring the spread of NDVI values.

Coefficient of variation (CoV): CoV can be used to compare the amount of variation in different sets of sample data. NDVI CoV images were generated by computing for each pixel the standard deviation (STD) of the set of individual NDVI values and dividing this by the mean (M) of these values. This represents the dispersion of NDVI values relative to the mean value.

Temporal trends: The long-term trends of the indicators of biological productivity may be taken as indicators of land degradation (where the trend is declining) or land improvement (where the trend is increasing). A positive change in the value of a pixel-level CoV over time relates to increased dispersion of values, not increasing NDVI; similarly, a negative CoV dispersion - which is the case over nearly the whole country - means decreasing dispersion of NDVI around mean values, not decreasing NDVI.

The patterns and trends of all NDVI indicators for each pixel, determined by the slope of the linear regression equation, are depicted in Figures S1-7; their values are summarised in Table S2. No further analyses were made for these indicators except for the sum NDVI which is discussed in detail in the main text. It is recommended, however, that these maps should be considered in the field investigation - in particular the land use change during the study period (1981-2003).




Figure S1. Annual minimum NDVI 1981-2003: Pattern (a), trends (b – percentage)




Figure S1. Annual minimum NDVI 1981-2003: Trend (c – absolute) and confidence levels (d)




Figure S2. Annual maximum NDVI 1981-2003: Pattern (a), trends (b – percentage)




Figure S2. Annual maximum NDVI 1981-2003: Pattern (a), trends (c – absolute) and confidence levels (d)




Figure S3. Annual max-min NDVI 1981-2003: Pattern (a), trends (b – percentage)




Figure S3. Annual max-min NDVI 1981-2003: trends (c – absolute) and confidence levels (d)




Figure S4. Annual mean NDVI 1981-2003: Pattern (a), trends (b – percentage)




Figure S4. Annual mean NDVI 1981-2003: trends (c – absolute) and confidence levels (d)




Figure S5. Annual sum NDVI 1981-2003: Pattern (a), trends (b – percentage)





Figure S5. Annual sum NDVI 1981-2003: trends (c – absolute) and confidence levels (d)

a



b




Figure S6. NDVI standard deviation 1981-2003: Pattern (a), trends (b – percentage)




Figure S6. NDVI standard deviation 1981-2003: Trends (c – absolute) and confidence levels (d)




Figure S7. NDVI coefficient of variation 1981-2003: Pattern (a), trends (b – percentage)




Figure S7. NDVI coefficient of variation 1981-2003: Trends (c – absolute) and confidence (d)



ISRIC - World Soil Information is an independent foundation with a global mandate, funded by the Netherlands Government. We have a strategic association with Wageningen University and Research Centre.

Our aims:

- *To inform and educate - through the World Soil Museum, public information, discussion and publication*
- *As ICSU World Data Centre for Soils, to serve the scientific community as custodian of global soil information*
- *To undertake applied research on land and water resources.*